

MRI-derived bone consistency index correlates to bone composition and mechanical stiffness

Abigail L. Honga , Mikayel Ispiryana , Mugdha V. Padalkarb , Brandon C. Jonesa,c , Alexandra S. Batzdorfa , Snehal S. Shetyec , urban center Pleshkob , Chamith S. Rajapaksea,c,

a Department of Radiology, University of Pennsylvania, us of America b Department of engineering science, Temple University, us of America c Department of orthopedical Surgery, University of Pennsylvania, us of America, E-mail: chamith@mail.med.upenn.edu

ABSTRACT

The MRI-derived consistency index (PI) could be a non-invasively obtained biomarker supported Associate in Nursing ultrashort echo time sequence that pictures each certain and pore water protons in bone, adore water guaranteed to organic scleroprotein matrix and freely moving water, severally. This live is understood to powerfully correlate with the particular meter animal tissue bone consistency. However, it's unknown whether or not PI might also be able to directly quantify bone organic composition and/or mechanical properties. we tend to investigated this in human body tibiae by scrutiny PI values to close infrared spectral imaging (NIRSI) integrative knowledge and mechanical compression knowledge. knowledge were obtained from a cohort of eighteen tibiae from male and feminine donors with a mean \pm Mount Rushmore State age of seventy \pm twenty one years. Biomechanical stiffness in compression and NIRSI-derived albuminoid and certain water content all had vital inverse correlations with PI ($r = -0.79, -0.73, \text{ and } -0.95$ and $p = \text{zero.002, 0.007, and } < 0.001$, respectively). The MRI-derived bone PI alone was a moderate predictor of bone stiffness ($R^2 = \text{zero.63, } p = 0.002$), and variable analyses showed that neither animal tissue bone cross-sectional space nor NIRSI values improved bone stiffness prediction compared to PI alone. However, NIRSI-obtained albuminoid and water

knowledge along were a moderate predictor of bone stiffness ($R^2 = \text{zero.52, } p = 0.04$). Our knowledge validates the MRI-derived consistency index as a powerful predictor of organic composition of bone and a moderate predictor of bone stiffness, and conjointly provides preliminary proof that NIRSI measures could also be helpful in future pre-clinical studies on bone pathology.

Bone fractures cause a high risk to the aging and unhealthy population, and assessments of bone mineral density (BMD) ar usually accustomed determine a patient's risk of fracture. for instance, various studies have shown that ladies with low bone density within the radius or bone ar at magnified risk of hip fracture, resonance imaging (MRI) ultrashort echo time (UTE) is a picture acquisition protocol that has incontestible hefty capability for imaging bone.

The borderline sample preparation, non-destructive nature of the scan, and relative speed of NIRSI makes it a perfect technique for investigation of changes in water content, distribution, and surroundings in pre-clinical studies of bone pathology and medical specialty.

Here, we tend to speculate that animal tissue PI will give correct measurements of bone organic material composition compared to NIRSI knowledge. we tend to additional speculate that,

thanks to its legendary ability to image structural knowledge of bone, PI can give correct predictions of bone strength.

Biomechanical testing leg bone segments twenty five metric linear unit long were obtained from the distal regions of the remains samples then underwent uniaxial compression tests employing a servo-hydraulic material testing machine (Instron

8874 Instron, Norwood, MA) equipped with a 100kN load cell.

The health and strength of bone is decided by various factors, together with the amount of mineralized tissue, the amount, distribution, and crosslinking ability of albuminoid, and therefore the macro- and micro-structure of bone, The bone is Associate in Nursing organic matrix mixture that's primarily comprised of albuminoid.

Keywords: MRI Porosity index Bone stiffness Near infrared spectral imaging Bone biomechanics Magnetic resonance imaging Ultrashort echo time