

## Finite element models for fracture prevention in patients with metastatic bone disease. A literature review

Amelie Sasa Esther Tanckb An Sermonc G. Harryvan Lenthe

Biomechanics Section, KU Leuven, Leuven, Belgium, E-mail: [harry.vanlenthe@kuleuven.be](mailto:harry.vanlenthe@kuleuven.be)

### ABSTRACT

Management of displaced intra-articular bone fracture has remained unclear for orthopedical surgeons and there's arguing to settle on the simplest strategies of treatment. wide selection of post-operative complications and sort of outcomes cause this complexity. Although, there's no sturdy proof, during which support superior advantage of bone affixation in treatment of DIACF, it's still a well-liked methodology. the target of this study was to gift outcome and complications of surgical management of DIACF with tri-cortical bone bone graft. cardinal cases containing forty seven men (57 feet) and 6 girls (7 feet) with DIACF underwent surgery by mean of tri-cortical machine graft bone from bone crest. AOFAS hind foot scale (consist of subjective and objective variables as well as 3 major classes pain, perform and alignment) and VAS questionnaires were used for full assessment of post-operative outcomes and complications. Mann-Whitney U check was used to compare means that. The mean  $\pm$  SD AOFAS hind foot score was eighty eight.  $18 \pm 7.12$ . AOFAS hind foot and VAS scores showed no important distinction between male and feminine. still, higher outcome of surgery among those below thirty five year mature supported AOFAS hind foot score. Use of bone graft for treatment of DIACF would possibly yield higher results, however, the results in all probability dissent in varied conditions and things. At the end, we have a tendency to suggest comparison results of various strategies of treatment of DIACF by one giant run with below the management of confounders. Calcaneal

fracture is that the commonest website of injury among tarsal bones and Intra-articular fractures accounted for regarding seventy fifth of all bone fractures. Displaced intra-articular bone fractures (DIACF) is sometimes created by vertical violence delivered to the foot. After a few years of studding, management of displaced intra-particular os tarsi fibulare fracture has remained unclear for orthopedical surgeons and there's arguing to settle on the simplest strategies of treatment like plates, bone graft and void filler. wide selection of post-operative complications and sort of outcomes cause this complexity. Wilmot administrated the primary use of bone graft for surgical management of DIACF. Although, there's no sturdy proof, during which support superior advantage of bone grafting in treatment of DIACF, it's still a well-liked methodology.

Study population of Associate in Nursing 100 and sixteen patients with calcaneal fracture were recruited for our study from Gregorian calendar month 1997 to could 2010 in Akhtar hospital, center of orthopedical surgery, Health Services and Medical Education, Tehran, Iran. Forty-five cases with extra-articular form of fracture were thought of to treat cautiously and were excluded from the study. Among seventy one eligible cases with displaced intra-articular bone fracture, those with but two years of follow up time and untouchable follow up knowledge were conjointly excluded. The operative procedure was performed once the soft tissue oedema attenuated and therefore the wrinkle

sign was positive on hind-foot's soft tissue in few days after trauma.

Conservative management of DIACF typically leads to complications like secondary inflammatory disease and malunion of the calcaneum. Therefore, ORIF is obtaining methodology of interest for treatment of DIACF. though the employment of bone grafting in management of DIACF have Associate in Nursing previous history in practice, its utility isn't still

clear. Rate of definitive and sometimes use of bone and sometimes ORIF has been reportable to be two hundredth and forty second in nationwide study in Kingdom of The Netherlands, severally [14]. Wang et al. used body covering reduction and internal fixation combined with fluid bone graft for treatment of fifteen cases of DIACF. After average of eighteen months of follow, they reportable no complication as well as wound infection, screw breakage and calcaneum varus.

**Keywords:** Bone metastases Fracture risk Prophylactic treatment Femoroplasty Finite element analysis